arXiv:1203.3250v2 [math.DG] 14 Nov 2012

REMARKS ON THE $\mathbb{R}/\mathbb{Z}$ GROTHENDIECK-RIEMANN-ROCH THEOREM

MAN-HO HO

Dedicated to my father Kar-Ming Ho

ABSTRACT. We give a direct proof that the $\mathbb{R}/\mathbb{Z}$ analytic index in $\mathbb{R}/\mathbb{Z}$ K-theory is well defined and the details of a short proof of the $\mathbb{R}/\mathbb{Z}$ Grothendieck-Riemann-Roch theorem stated in a paper of Bismut. The proofs of these theorem do not make use differential K-theory.

CONTENTS

1. Introduction 1
2. Differential characters 3
2.1. Multiplication of differential characters 4
2.2. Pushforward of differential characters 4
3. The $\mathbb{R}/\mathbb{Z}$ K-theory 5
3.1. The $\mathbb{R}/\mathbb{Z}$ K-group 5
3.2. The $\mathbb{R}/\mathbb{Z}$ analytic index 6
4. Main results 8
4.1. The $\mathbb{R}/\mathbb{Z}$ analytic index is well defined 8
4.2. The $\mathbb{R}/\mathbb{Z}$ Grothendieck-Riemann-Roch theorem 10
References 12

1. INTRODUCTION

We give a direct proof that the $\mathbb{R}/\mathbb{Z}$ analytic index

$$\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}} : K^{-1}_{\mathrm{L}}(X; \mathbb{R}/\mathbb{Z}) \to K^{-1}_{\mathrm{L}}(B; \mathbb{R}/\mathbb{Z})$$

[11, Definition 14] is well defined, and the details of a short proof of the $\mathbb{R}/\mathbb{Z}$ Grothendieck-Riemann-Roch theorem [11, Corollary 4]

$$\begin{tikzcd} K^{-1}_{\mathrm{L}}(X; \mathbb{R}/\mathbb{Z}) \arrow[r, "\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}"] \arrow[d, "\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}"'] & H^{\mathrm{odd}}(B; \mathbb{R}/\mathbb{Q}) \arrow[d, "\int_{X/B} \mathrm{Todd}(X/B) \cup (\cdot)"] \\ K^{-1}_{\mathrm{L}}(B; \mathbb{R}/\mathbb{Z}) \arrow[r, "\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}"'] & H^{\mathrm{odd}}(B; \mathbb{R}/\mathbb{Q}) \end{tikzcd} \tag{1}$$

2010 *Mathematics Subject Classification.* Primary 19K56, 58J20, 53C08, 19L10.

stated in [3, p.17], without appealing to differential K-theory.

Let $\pi : X \to B$ be a proper submersion of manifolds of even relative dimension with closed Spin^c-fibers. The cohomological version of the Atiyah-Singer family index theorem, i.e., the Grothendieck-Riemann-Roch theorem, expresses the Chern character of the index bundle in topological means:

$$\mathrm{ch}(\mathrm{ind}^{\mathrm{an}}(E)) = \int_{X/B} \mathrm{Todd}(X/B) \cup \mathrm{ch}(E). \tag{2}$$

The Atiyah-Singer family index theorem is an integral theorem, i.e., $K^0(*) = \mathbb{Z}$. From (2) we see that if $\mathrm{ch}(E) = 0$, then $\mathrm{ch}(\mathrm{ind}^{\mathrm{an}}(E)) = 0$. As explained in [11], this is the motivation of having a geometric model for the $\mathbb{R}/\mathbb{Z}$ K-group $K^{-1}(X; \mathbb{R}/\mathbb{Z})$ [1] and proving an index theorem taking values in $\mathbb{R}/\mathbb{Z}$ for virtual bundles with vanishing Chern character.

The $\mathbb{R}/\mathbb{Z}$ family index theorem $\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}} = \mathrm{ind}^{\mathrm{topo}}_{\mathrm{L}}$ [11, Corollary 3], equates the $\mathbb{R}/\mathbb{Z}$ analytic index $\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}$ with the $\mathbb{R}/\mathbb{Z}$ topological index $\mathrm{ind}^{\mathrm{topo}}_{\mathrm{L}}$ [11, §4]. The $\mathbb{R}/\mathbb{Z}$ Grothendieck-Riemann-Roch theorem (1) is obtained by applying the $\mathbb{R}/\mathbb{Z}$ Chern character

$$\mathrm{ch}_{\mathbb{R}/\mathbb{Q}} : K^{-1}_{\mathrm{L}}(X; \mathbb{R}/\mathbb{Z}) \to H^{\mathrm{odd}}(X; \mathbb{R}/\mathbb{Q})$$

(also known as the Cheeger-Chern-Simons class) to the $\mathbb{R}/\mathbb{Z}$ family index theorem.

On the other hand, $K^{-1}_{\mathrm{L}}(X; \mathbb{R}/\mathbb{Z})$ is now known as the flat part of differential K-theory $\widehat{K}_{\mathrm{FL}}(X)$ [7]. Thus we can consider the $\mathbb{R}/\mathbb{Z}$ analytic index $\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}$ as a special case [7, Proposition 7.37] of the differential analytic index $\mathrm{ind}^{\mathrm{an}}_{\mathrm{FL}}$ [7, Definition 7.27], and the $\mathbb{R}/\mathbb{Z}$ topological index as a special case [7, Proposition 8.10] of the Freed-Lott differential topological index $\mathrm{ind}^{\mathrm{topo}}_{\mathrm{FL}}$ [7, Definition 5.34]. Therefore the $\mathbb{R}/\mathbb{Z}$ family index theorem is a special case of the differential family index theorem

$$\mathrm{ind}^{\mathrm{an}}_{\mathrm{FL}} = \mathrm{ind}^{\mathrm{topo}}_{\mathrm{FL}} : \widehat{K}_{\mathrm{FL}}(X)\widehat{K}_{\mathrm{FL}}(B)$$

[7, Theorem 7.35]; and the $\mathbb{R}/\mathbb{Z}$ Grothendieck-Riemann-Roch theorem is a special case of the differential Grothendieck-Riemann-Roch theorem [7, Corollary 8.26]

$$\widehat{\mathrm{ch}}_{\mathrm{FL}}(\mathrm{ind}^{\mathrm{an}}_{\mathrm{FL}}(\mathcal{E})) = \widehat{\int_{X/B}} \widehat{\mathrm{Todd}}(T^V X, \widehat{\nabla}^{T^V X}) * \widehat{\mathrm{ch}}_{\mathrm{FL}}(\mathcal{E}) \in \widehat{H}^{\mathrm{even}}(B; \mathbb{R}/\mathbb{Q}),$$

since $H^{\mathrm{odd}}(B; \mathbb{R}/\mathbb{Q})$ embeds in $\widehat{H}^{\mathrm{even}}(B; \mathbb{R}/\mathbb{Q})$, where $\widehat{H}^{\mathrm{even}}(B; \mathbb{R}/\mathbb{Q})$ is the ring of Cheeger-Simons differential characters.

Thus the direct proof of the differential analytic index $\mathrm{ind}^{\mathrm{an}}_{\mathrm{FL}}$ is well defined [8, Proposition 3] and the condensed proof of the differential Grothendieck-Riemann-Roch theorem [8, Theorem 1] imply a direct proof of the $\mathbb{R}/\mathbb{Z}$ analytic index $\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}$ is well defined and a short proof of the $\mathbb{R}/\mathbb{Z}$ Grothendieck-Riemann-Roch theorem.

The purpose of this paper is to give a direct proof that the $\mathbb{R}/\mathbb{Z}$ analytic index $\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}$ is well defined (Proposition 3), and the details of a short proof of the $\mathbb{R}/\mathbb{Z}$ Grothendieck-Riemann-Roch theorem as stated in [3, p. 17]. These proofs do not make use of differential K-theory.

The paper is organized as follows: Section 2 contains all the necessary background materials we need, including Cheeger-Simons differential characters, their multiplication and some properties of pushforward; the $\mathbb{R}/\mathbb{Z}$ K-group $K_{\mathrm{L}}^{-1}(X;\mathbb{R}/\mathbb{Z})$ and the construction of $\mathbb{R}/\mathbb{Z}$ analytic index $\mathrm{ind}_{\mathrm{L}}^{\mathrm{an}}$. In Section 3 we prove the main results of this paper.

2. Differential characters

We recall Cheeger–Simons differential characters [5] with coefficients in $\mathbb{R}/\mathbb{Q}$. Let X be a manifold. The ring of differential characters of degree $k \geq 1$ is

$$\widehat{H}^k(X;\mathbb{R}/\mathbb{Q}) = \{f \in \mathrm{Hom}(Z_{k-1}(X),\mathbb{R}/\mathbb{Q}) | \exists \omega_f \in \Omega^k(X) \text{ such that } f \circ \partial = \overline{\omega_f}\},$$

where $^{-} : \Omega^k(X) \to C^k(X;\mathbb{R}/\mathbb{Q})$ is an injective homomorphism defined by $\overline{\omega}(c_k) := \displaystyle\int_{c_k} \omega \mod \mathbb{Q}$. It is easy to show that ω_f is a closed k-form with periods in $\mathbb{Q}$ and is uniquely determined by $f \in \widehat{H}^k(X;\mathbb{R}/\mathbb{Q})$. In the following hexagon, the diagonal sequences are exact, and every triangle and square commutes [5, Theorem 1.1]:

$$\begin{tikzcd}
0 \arrow[dr] & & & & 0 \\
& H^{k-1}(X;\mathbb{R}/\mathbb{Q}) \arrow[rr, "-B"] \arrow[dr, "i_1"] & & H^k(X;\mathbb{Q}) \arrow[ur] \arrow[dr, "r"] & \\
H^{k-1}(X;\mathbb{R}) \arrow[ur, "\alpha"] \arrow[dr, "\beta"'] & & \widehat{H}^k(X;\mathbb{R}/\mathbb{Q}) \arrow[ur, "\delta_2"] \arrow[dr, "\delta_1"] & & H^k(X;\mathbb{R}) \\
& \frac{\Omega^{k-1}(X)}{\Omega^{k-1}_{\mathbb{Q}}(X)} \arrow[rr, "d"'] \arrow[ur, "i_2"] & & \Omega^k_{\mathbb{Q}}(X) \arrow[ur, "s"] \arrow[dr] & \\
0 \arrow[ur] & & & & 0
\end{tikzcd} \tag{3}$$

The maps are defined as follows: r is induced by $\mathbb{Q} \hookrightarrow \mathbb{R}$,

$$i_1([z]) = z|_{Z_{k-1}(X)},\ i_2(\omega) = \overline{\omega}|_{Z_{k-1}(X)},\ \delta_1(f) = \omega_f \text{ and } \delta_2(f) = [c],$$

where $[c] \in H^k(X;\mathbb{Q})$ is the unique cohomology class satisfying $r[c] = [\omega_f]$, and $\Omega^k_{\mathbb{Q}}(X)$ consists of closed forms with periods in $\mathbb{Q}$. (We will not use the other maps.)

Invariant polynomials for $\mathrm{U}(n)$ have associated characteristic classes and differential characters. In particular, for a Hermitian vector bundle $E \to X$

with a metric h and a unitary connection ∇, the differential Chern character is the unique natural differential character [5, Theorem 2.2]

$$\widehat{\mathrm{ch}}(E, h, \nabla) \in \widehat{H}^{\mathrm{even}}(X; \mathbb{R}/\mathbb{Q}) \tag{4}$$

such that

$$\delta_1(\widehat{\mathrm{ch}}(E, h, \nabla)) = \mathrm{ch}(\nabla) \text{ and } \delta_2(\widehat{\mathrm{ch}}(E, h, \nabla)) = \mathrm{ch}(E).$$

We will write $\widehat{\mathrm{ch}}(E, h, \nabla)$ as $\widehat{\mathrm{ch}}(E, \nabla)$ in the sequel.

2.1. **Multiplication of differential characters.** In [5] multiplication of differential characters is defined. Let $E : \Omega^{k_1}(X) \times \Omega^{k_2}(X) \to C^{k_1+k_2-1}(X; \mathbb{R})$ be a natural chain homotopy between the wedge product $\wedge$ and the cup product $\cup$, i.e., for $\omega_i \in \Omega^{k_i}(X)$, we have

$$\delta E(\omega_1, \omega_2) + E(d\omega_1, \omega_2) + (-1)^{k_1} E(\omega_1, d\omega_2) = \omega_1 \wedge \omega_2 - \omega_1 \cup \omega_2$$

as cochains. Note that any two choices of E are naturally chain homotopic. For $f \in \widehat{H}^{k_1}(X; \mathbb{R}/\mathbb{Q})$ and $g \in \widehat{H}^{k_2}(X; \mathbb{R}/\mathbb{Q})$, define $f * g \in \widehat{H}^{k_1+k_2}(X; \mathbb{R}/\mathbb{Q})$ by

$$f * g = \big(\widetilde{T_f \cup \omega_g} + (-1)^{k_1} \widetilde{\omega_f \cup T_g} + \widetilde{T_f \cup \delta T_g} + E\widetilde{(\omega_f, \omega_g)}\big)|_{Z_{k_1+k_2-1}(X)},$$

where $T_f, T_g \in C^{k-1}(X, \mathbb{R})$ are lifts of f and g.

Proposition 1. [5, Theorem 1.11] Let $f \in \widehat{H}^{k_1}(X; \mathbb{R}/\mathbb{Q})$ and $g \in \widehat{H}^{k_2}(X; \mathbb{R}/\mathbb{Q})$.

1. $f * g$ is independent of the choice of the lifts T_f and T_g,
2. $f * (g * h) = (f * g) * h$ and $f * g = (-1)^{k_1 k_2} g * f$,
3. $\omega_{f*g} = \omega_f \wedge \omega_g$ and $c_{f*g} = c_f \cup c_g$. i.e., δ_1 and δ_2 are ring homomorphisms,
4. If $\phi : N \to M$ is a smooth map, then $\phi^*(f * g) = \phi^*(f) * \phi^*(g)$,
5. If $\theta \in \Omega^\bullet(X)$, then $i_2(\theta) * f = i_2(\theta \wedge \omega_f)$,
6. If $[c] \in H^\bullet(X; \mathbb{R}/\mathbb{Q})$, then $f * i_1([c]) = (-1)^{k_1} i_1([c_f] \cup [c])$.

2.2. **Pushforward of differential characters.** The pushforward of differential characters is defined in [9, §3.4]. We only consider proper submersions $\pi : X \to B$ with closed fibers of relative dimension n, where the definition [7, §8.3] is straightforward: for $k \geq n$,

$$\widehat{\int_{X/B}} : \widehat{H}^k(X; \mathbb{R}/\mathbb{Q}) \to \widehat{H}^{k-n}(B; \mathbb{R}/\mathbb{Q}), \ \left(\widehat{\int_{X/B}} f\right)(z) = f(\pi^{-1}(z)).$$

Let $\displaystyle\int_{X/B}$ denote both the pushforward of forms and cohomology classes.

Proposition 2. [9, §3.4] Let $f \in \widehat{H}^k(X; \mathbb{R}/\mathbb{Q})$, $[c] \in H^{k-1}(X; \mathbb{R}/\mathbb{Q})$ and $\theta \in \dfrac{\Omega^{k-1}(X)}{\Omega^{k-1}_{\mathbb{Q}}(X)}$. Then

(1) $\delta_1\Big(\widehat{\int_{X/B}} f\Big) = \int_{X/B} \omega_f.$

(2) $\delta_2\Big(\widehat{\int_{X/B}} f\Big) = \int_{X/B} [c_f].$

(3) $\widehat{\int_{X/B}} i_1([c]) = i_1\Big(\int_{X/B} [c]\Big).$

(4) $\widehat{\int_{X/B}} i_2(\theta) = i_2\Big(\int_{X/B} \theta\Big).$

3. The $\mathbb{R}/\mathbb{Z}$ K-theory

3.1. **The $\mathbb{R}/\mathbb{Z}$ K-group.** In this subsection we recall the definition of K_{L}^{-1}.

Let $E \to X$ be a Hermitian vector bundle with a Hermitian metric h, ∇ a unitary connection. Let $\omega \in \dfrac{\Omega^{\mathrm{odd}}(X)}{\mathrm{Im}(d)}$. $K_{\mathrm{L}}^{-1}(X;\mathbb{R}/\mathbb{Z})$ [11, Definition 7] is the abelian group given by generators $\mathcal{E} = (E, h, \nabla, \omega)$ such that $\mathrm{ch}(\nabla) - \mathrm{rank}(E) = -d\omega$ and two relations:

(1) $\mathcal{E}_2 = \mathcal{E}_1 + \mathcal{E}_3$ in $K_{\mathrm{L}}^{-1}(X;\mathbb{R}/\mathbb{Z})$ if and only if there exists a short exact sequence of vector bundles
$$0 \longrightarrow E_1 \longrightarrow E_2 \longrightarrow E_3 \longrightarrow 0$$
and $\omega_2 = \omega_1 + \omega_3 + \mathrm{CS}(\nabla^1, \nabla^2, \nabla^3)$.

(2) The virtual rank of any arbitrary element in $K^{-1}(X;\mathbb{R}/\mathbb{Z})$ is zero.

In particular, $\mathcal{E}_1 = \mathcal{E}_2 \in K_{\mathrm{L}}^{-1}(X;\mathbb{R}/\mathbb{Z})$ if and only if there exists a generator (G, h^G, ∇^G, ϕ) such that $E_1 \oplus G \cong E_2 \oplus G$ and $\omega_2 - \omega_1 = \mathrm{CS}(\nabla^1 \oplus \nabla^G, \nabla^2 \oplus \nabla^G)$.

By a $\mathbb{Z}_2$-graded element $\mathcal{E} = (E, h^E, \nabla^E, \phi) \in K_{\mathrm{L}}^{-1}(X;\mathbb{R}/\mathbb{Z})$ we mean $E = E^+ \oplus E^-$ is a $\mathbb{Z}_2$-graded complex vector bundle over X, $h^E = h^+ \oplus h^-$ and $\nabla^E = \nabla^+ \oplus \nabla^-$, where $h^\pm$ is a Hermitian metric and $\nabla^\pm$ is a unitary connection on $E^\pm \to X$, and $\mathrm{ch}(\nabla^E) := \mathrm{ch}(\nabla^+) - \mathrm{ch}(\nabla^-) = -d\phi$. Given an element in $K_{\mathrm{L}}^{-1}(X;\mathbb{R}/\mathbb{Z})$, we can define a $\mathbb{Z}_2$-graded element in $K_{\mathrm{L}}^{-1}(X;\mathbb{R}/\mathbb{Z})$, and vice versa [11, p. 8].

The following sequence is exact [10, §7.21] (see also [11, (13)])
$$\begin{aligned} K^{-1}(X) \xrightarrow{r\mathrm{ch}^{\mathrm{odd}}} H^{\mathrm{odd}}(X;\mathbb{R}) &\xrightarrow{\ \alpha\ } K_{\mathrm{L}}^{-1}(X;\mathbb{R}/\mathbb{Z}) \xrightarrow{\ \beta\ } \\ K(X) &\xrightarrow{r\mathrm{ch}} H^{\mathrm{even}}(X;\mathbb{R}) \end{aligned} \tag{5}$$
where r is induced by the inclusion of coefficients $\mathbb{Q} \hookrightarrow \mathbb{R}$, and
$$\alpha([\omega]) = (\mathbb{C}^n, h, \nabla^{\mathrm{flat}}, \omega) - (\mathbb{C}^n, h, \nabla^{\mathrm{flat}}, 0),$$
$$\beta((E, h^E, \nabla^E, \phi^E) - (F, h^F, \nabla^F, \phi^F)) = [E] - [F].$$

The $\mathbb{R}/\mathbb{Z}$ Chern character $\mathrm{ch}_{\mathbb{R}/\mathbb{Q}} : K_{\mathrm{L}}^{-1}(X;\mathbb{R}/\mathbb{Z}) \to H^{\mathrm{odd}}(X;\mathbb{R}/\mathbb{Q})$ [11, Definition 9] is defined by

$$\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathcal{E}) = \left[\frac{1}{k}\,\mathrm{CS}(k\nabla^{E}, \nabla_0^{kE}) + \omega\right] \mod \mathbb{Q}, \tag{6}$$

where $k \in \mathbb{N}$ and ∇_0^{kE} is a unitary connection on $kE \cong \mathbb{C}^N$, for some $N \in \mathbb{N}$, with trivial holonomy. [11, Lemma 1, Proposition 1] show that $\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}$ is well defined. i.e., $\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}$ is independent of the choices of k and ∇_0^{kE}.

The $\mathbb{R}/\mathbb{Z}$ Chern character of a $\mathbb{Z}_2$-graded element $\mathcal{E}$ is given by

$$\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathcal{E}) = \left[\frac{1}{k}\,\mathrm{CS}(k\nabla^{+}, j^*k\nabla^{-}) + \omega\right] \mod \mathbb{Q}, \tag{7}$$

where $j : kE^+ \to kE^-$ is an isometric isomorphism. It is independent of the choices of k and j [11, p. 12].

Note that the following diagram commutes.

$$\begin{tikzcd} H^{\mathrm{odd}}(X;\mathbb{R}) \arrow[r, "\alpha"] \arrow[dr, "\alpha'"'] & K_{\mathrm{L}}^{-1}(X;\mathbb{R}/\mathbb{Z}) \arrow[d, "\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}"] \\ & H^{\mathrm{odd}}(X;\mathbb{R}/\mathbb{Q}) \end{tikzcd} \tag{8}$$

where α' is given in (4). By the definition of α, we have

$$\begin{aligned} \mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\alpha([\omega])) &= \mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathbb{C}^n, h, \nabla^{\mathrm{flat}}, \omega) - \mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathbb{C}^n, h, \nabla^{\mathrm{flat}}, 0) \\ &= [\mathrm{CS}(\nabla^{\mathrm{flat}}, \nabla_0^{\mathbb{C}^n}) + \omega] - [\mathrm{CS}(\nabla^{\mathrm{flat}}, \nabla_0^{\mathbb{C}^n})] \mod \mathbb{Q} \\ &= [\omega] \mod \mathbb{Q} \\ &= \alpha'([\omega]). \end{aligned}$$

3.2. **The $\mathbb{R}/\mathbb{Z}$ analytic index.** The $\mathbb{R}/\mathbb{Z}$ analytic index of a $\mathbb{Z}_2$-graded generator $\mathcal{E} = (E, h, \nabla, \phi) \in K_{\mathrm{L}}^{-1}(X;\mathbb{R}/\mathbb{Z})$ is roughly given by the geometric construction of the analytic index of (E, h, ∇) with a modified pushforward of the form ϕ.

First of all we recall the Bismut's construction of the analytic index. Let $\pi : X \to B$ be a proper submersion of even relative dimension n, and let $T^VX \to X$ be the vertical tangent bundle, which is assumed to have a metric g^{T^VX}. A given horizontal distribution $T^HX \to X$ and a Riemannian metric g^{TB} on B determine a metric on $TX \to X$ by $g^{TX} := g^{T^VX} \oplus \pi^* g^{TB}$. If ∇^{TX} is the corresponding Levi-Civita connection, then $\nabla^{T^VX} := P \circ \nabla^{TX} \circ P$ is a connection on $T^VX \to X$, where $P : TX \to T^VX$ is the orthogonal projection. $T^VX \to X$ is assumed to have a Spin^c structure. Denote by $S^VX \to X$ the spinor^c bundle associated to the characteristic Hermitian line bundle $L^VX \to X$ with a unitary connection ∇^{L^VX}. Define a connection $\widehat{\nabla}^{T^VX}$ on $S^VX \to X$ by $\widehat{\nabla}^{T^VX} := \nabla^{T^VX} \otimes \nabla^{L^V}$, where ∇^{T^VX} also denotes the lift of ∇^{T^VX} to the local spinor bundle. The Todd form $\mathrm{Todd}(\widehat{\nabla}^{T^VX})$ of S^VX is related to the $\hat{A}$-form of T^VX by

$$\mathrm{Todd}(\widehat{\nabla}^{T^VX}) = \widehat{A}(\nabla^{T^VX}) \wedge e^{\frac{1}{2}c_1(\nabla^{L^VX})}.$$

For $k \geq n$, the modified pushforward of forms $\pi_* : \Omega^k(X) \to \Omega^{k-n}(B)$ [7, (3.2)] is defined by

$$\pi_*(\phi) = \int_{X/B} \mathrm{Todd}(\widehat{\nabla}^{T^VX}) \wedge \phi.$$

It induces a map, still denoted by $\pi_* : \dfrac{\Omega^{\mathrm{odd}}(X)}{\mathrm{Im}(d)} \to \dfrac{\Omega^{\mathrm{odd}}(B)}{\mathrm{Im}(d)}$.

We briefly recall the definition of the Bismut–Cheeger eta form $\widetilde{\eta}(\mathcal{E}) \in \dfrac{\Omega^{\mathrm{odd}}(B)}{\mathrm{Im}(d)}$ associated to $\mathcal{E} \in \widehat{K}_{\mathrm{FL}}(X)$. With the above setup, consider the infinite-rank superbundle $\pi_*E \to B$, where the fibers at each $b \in B$ is given by

$$(\pi_*E)_b := \Gamma(X_b, (S^VX \otimes E)|_{X_b}).$$

Recall that $\pi_*E \to B$ admits an induced Hermitian metric and a connection ∇^{π_*E} compatible with the metric [2, §9.2, Proposition 9.13]. For each $b \in B$, the canonically constructed Dirac operator

$$\mathsf{D}^E_b : \Gamma(X_b, (S^VX \otimes E)|_{X_b}) \to \Gamma(X_b, (S^VX \otimes E)|_{X_b})$$

gives a family of Dirac operators, denoted by $\mathsf{D}^E : \Gamma(X, S^VX \otimes E) \to \Gamma(X, S^VX \otimes E)$. Assume the family of kernels $\ker(\mathsf{D}^E_b)$ has locally constant dimension, i.e., $\ker(\mathsf{D}^E) \to B$ is a finite-rank Hermitian superbundle. Let $P : \pi_*E \to \ker(\mathsf{D}^E)$ be the orthogonal projection, $h^{\ker(\mathsf{D}^E)}$ be the Hermitian metric on $\ker(\mathsf{D}^E) \to B$ induced by P, and $\nabla^{\ker(\mathsf{D}^E)} := P \circ \nabla^{\pi_*E} \circ P$ be the connection on $\ker(\mathsf{D}^E) \to B$ compatible to $h^{\ker(\mathsf{D}^E)}$.

The (scaled) Bismut-superconnection $\mathbb{A}_t : \Omega(B, \pi_*E) \to \Omega(B, \pi_*E)$ [**?**, Definition 3.2] (see also [2, Proposition 10.15] and [6, (1.4)]), is defined by

$$\mathbb{A}_t := \sqrt{t}\mathsf{D}^E + \nabla^{\pi_*E} - \frac{c(T)}{4\sqrt{t}},$$

where $c(T)$ is the Clifford multiplication by the curvature 2-form of the fiber bundle. The Bismut–Cheeger eta form $\widetilde{\eta}(\mathcal{E})$ [4, (2.26)] (see also [6] and [2, Theorem 10.32]) is defined by

$$\widetilde{\eta}(\mathcal{E}) := \frac{1}{2\sqrt{\pi}} \int_0^\infty \frac{1}{\sqrt{t}} \mathrm{Str}\left(\frac{d\mathbb{A}_t}{dt} e^{-\mathbb{A}_t^2}\right) dt.$$

It satisfies

$$d\widetilde{\eta}(\mathcal{E}) = \int_{X/B} \mathrm{Todd}(\widehat{\nabla}^{T^VX}) \wedge \mathrm{ch}(\nabla^E) - \mathrm{ch}(\nabla^{\ker(\mathsf{D}^E)}).$$

Now we recall the definition of $\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}} : K^{-1}_{\mathrm{L}}(X;\mathbb{R}/\mathbb{Z}) \to K^{-1}_{\mathrm{L}}(B;\mathbb{R}/\mathbb{Z})$. Let $(E, h^E, \nabla^E, \phi) \in K^{-1}_{\mathrm{L}}(X;\mathbb{R}/\mathbb{Z})$ be a $\mathbb{Z}_2$-graded element. Since $E = E^+ \oplus E^-$ is a $\mathbb{Z}_2$-graded complex vector bundle, $h = h^+ \oplus h^-$ is a $\mathbb{Z}_2$-graded Hermitian metric and $\nabla = \nabla^+ \oplus \nabla^-$ is a $\mathbb{Z}_2$-graded unitary connection on $E = E^+ \oplus E^-$, we apply the above geometric construction of the index bundle to each of $E^+ \to B$ and $E^- \to B$. Since each of the index bundle $\ker(\mathsf{D}^{E^\pm}) \to B$ is a $\mathbb{Z}_2$-graded bundle

$$\begin{aligned}\ker(\mathsf{D}^{E^+}) &= \ker(\mathsf{D}^{E^+})^+ \oplus \ker(\mathsf{D}^{E^+})^-,\\ \ker(\mathsf{D}^{E^-}) &= \ker(\mathsf{D}^{E^-})^+ \oplus \ker(\mathsf{D}^{E^-})^-,\end{aligned}$$

it follows that the index bundle of $E \to X$ is $\mathbb{Z}_2$-graded complex vector bundle defined by

$$\mathrm{ind}^{\mathrm{an}}(E) = \mathrm{ind}^{\mathrm{an}}(E)^+ \oplus \mathrm{ind}^{\mathrm{an}}(E)^-,$$

where

$$\begin{aligned}\mathrm{ind}^{\mathrm{an}}(E)^+ &= \ker(\mathsf{D}^{E^+})^+ \oplus \ker(\mathsf{D}^{E^-})^-\\ \mathrm{ind}^{\mathrm{an}}(E)^- &= \ker(\mathsf{D}^{E^+})^- \oplus \ker(\mathsf{D}^{E^-})^+.\end{aligned}$$

$\mathrm{ind}^{\mathrm{an}}(E) \to B$ inherits a L^2-metric $h^{\mathrm{ind}^{\mathrm{an}}(E)} = h^{\mathrm{ind}^{\mathrm{an}}(E)^+} \oplus h^{\mathrm{ind}^{\mathrm{an}}(E)^-}$, and a metric superconnection $\nabla^{\mathrm{ind}^{\mathrm{an}}(E)} = \nabla^{\mathrm{ind}^{\mathrm{an}}(E)^+} \oplus \nabla^{\mathrm{ind}^{\mathrm{an}}(E)^-}$.

The $\mathbb{R}/\mathbb{Z}$ analytic index $\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}} : K^{-1}_{\mathrm{L}}(X;\mathbb{R}/\mathbb{Z}) \to K^{-1}_{\mathrm{L}}(B;\mathbb{R}/\mathbb{Z})$ for a $\mathbb{Z}_2$-graded element $\mathcal{E} = (E, h, \nabla, \phi)$ is defined by

$$\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}(\mathcal{E}) := \left(\mathrm{ind}^{\mathrm{an}}(E), h^{\mathrm{ind}^{\mathrm{an}}(E)}, \nabla^{\mathrm{ind}^{\mathrm{an}}(E)}, \int_{X/B} \mathrm{Todd}(\widehat{\nabla}^{T^V X}) \wedge \omega + \widetilde{\eta}(\mathcal{E})\right).$$

If we consider a $\mathbb{Z}_2$-graded element $\mathcal{E} = (E, h, \nabla)$ to be $\mathcal{E} = \mathcal{E}^+ \oplus \mathcal{E}^-$, then

$$\widetilde{\eta}(\mathcal{E}) = \widetilde{\eta}(\mathcal{E}^+) + \widetilde{\eta}(\mathcal{E}^-).$$

4. MAIN RESULTS

4.1. **The $\mathbb{R}/\mathbb{Z}$ analytic index is well defined.** In this subsection we give a direct proof that the $\mathbb{R}/\mathbb{Z}$ analytic index $\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}$ is well defined.

Proposition 3. Let $\pi : X \to B$ be a proper submersion with closed Spin^c fibers of even relative dimension. The $\mathbb{R}/\mathbb{Z}$ analytic index

$$\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}} : K^{-1}_{\mathrm{L}}(X;\mathbb{R}/\mathbb{Z}) \to K^{-1}_{\mathrm{L}}(B;\mathbb{R}/\mathbb{Z})$$

is well defined.

Proof. Suppose $\mathcal{E}_i = (E_i, h^i, \nabla^i, \phi^i) \in K^{-1}_{\mathrm{L}}(X;\mathbb{R}/\mathbb{Z})$, for $i = 1, 2$, are $\mathbb{Z}_2$-graded elements such that $\mathcal{E}_1 - \mathcal{E}_2 = 0$. We prove that $\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}(\mathcal{E}_1) - \mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}(\mathcal{E}_2) = 0$ in $K^{-1}(B;\mathbb{R}/\mathbb{Z})$. Since $\mathcal{E}_1 = \mathcal{E}_2$, there exists a generator $(G, h^G, \nabla^G, \phi^G)$ such that

$$\begin{aligned}&E_1 \oplus G \cong E_2 \oplus G\\ &\phi^2 - \phi^1 = \mathrm{CS}(\nabla^1 \oplus \nabla^G, \nabla^2 \oplus \nabla^G)\end{aligned} \tag{9}$$

Since the index bundle is additive, it follows that

$$\begin{aligned}\operatorname{ind}^{\mathrm{an}}(E_1)\oplus\operatorname{ind}^{\mathrm{an}}(G) &\cong \operatorname{ind}^{\mathrm{an}}(E_1\oplus G)\\ &\cong \operatorname{ind}^{\mathrm{an}}(E_2\oplus G)\\ &\cong \operatorname{ind}^{\mathrm{an}}(E_2)\oplus\operatorname{ind}^{\mathrm{an}}(G)\end{aligned}\tag{10}$$

The commutativity of the diagram

$$\begin{tikzcd} K^{-1}(X;\mathbb{R}/\mathbb{Z}) \arrow[r, "\beta"] \arrow[d, "\operatorname{ind}^{\mathrm{an}}_{\mathrm{L}}"'] & K(X) \arrow[d, "\operatorname{ind}^{\mathrm{an}}"] \\ K^{-1}(B;\mathbb{R}/\mathbb{Z}) \arrow[r, "\beta"'] & K(B) \end{tikzcd}$$

and (10) implies that

$$\beta(\operatorname{ind}^{\mathrm{an}}_{\mathrm{L}}(\mathcal{E}_1)-\operatorname{ind}^{\mathrm{an}}_{\mathrm{L}}(\mathcal{E}_2))=\operatorname{ind}^{\mathrm{an}}(E_1)-\operatorname{ind}^{\mathrm{an}}(E_2)=0.$$

It follows from the exact sequence (5) that there exists $[\omega]\in H^{\mathrm{odd}}(B;\mathbb{R})$ such that $\alpha([\omega])=\operatorname{ind}^{\mathrm{an}}(\mathcal{E}_1)-\operatorname{ind}^{\mathrm{an}}(\mathcal{E}_2)$, i.e.,

$$\begin{aligned}&\Big(\operatorname{ind}^{\mathrm{an}}(E_1),h^{\operatorname{ind}^{\mathrm{an}}(E_1)},\nabla^{\operatorname{ind}^{\mathrm{an}}(E_1)},\int_{X/B}\operatorname{Todd}(\widehat{\nabla}^{T^VX})\wedge\phi^1+\widetilde{\eta}(\mathcal{E}_1)\Big)\\ &\qquad-\Big(\operatorname{ind}^{\mathrm{an}}(E_2),h^{\operatorname{ind}^{\mathrm{an}}(E_2)},\nabla^{\operatorname{ind}^{\mathrm{an}}(E_2)},\int_{X/B}\operatorname{Todd}(\widehat{\nabla}^{T^VX})+\widetilde{\eta}(\mathcal{E}_2)\Big)\\ &=(\mathbb{C}^n,h,\nabla^{\mathrm{flat}},\omega)-(\mathbb{C}^n,h,\nabla^{\mathrm{flat}},0).\end{aligned}$$

Thus, by definition, we have

$$\begin{aligned}&\omega+\int_{X/B}\operatorname{Todd}(\widehat{\nabla}^{T^VX})\wedge\phi^2+\widetilde{\eta}(\mathcal{E}_2)-\int_{X/B}\operatorname{Todd}(\widehat{\nabla}^{T^VX})\wedge\phi^1-\widetilde{\eta}(\mathcal{E}_1)\\ &=\operatorname{CS}(\nabla^{\operatorname{ind}^{\mathrm{an}}(E_1)}\oplus\nabla^{\mathrm{flat}}\oplus,\nabla^{\operatorname{ind}^{\mathrm{an}}(E_2)}\oplus\nabla^{\mathrm{flat}})\quad\operatorname{mod}\ \Omega^{\mathrm{odd}}_{\mathrm{exact}}(B).\end{aligned}\tag{11}$$

By (9) and (10) we have

$$\begin{aligned}\phi^2-\phi^1&=\operatorname{CS}(\nabla^1\oplus\nabla^G,\nabla^2\oplus\nabla^G),\\ \operatorname{ind}^{\mathrm{an}}(E_1)\oplus\operatorname{ind}^{\mathrm{an}}(G)&\cong\operatorname{ind}^{\mathrm{an}}(E_2)\oplus\operatorname{ind}^{\mathrm{an}}(G),\end{aligned}$$

it follows that (11) becomes

$$\begin{aligned}&\omega+\int_{X/B}\operatorname{Todd}(\widehat{\nabla}^{T^VX})\wedge\operatorname{CS}(\nabla^1\oplus\nabla^G,\nabla^2\oplus\nabla^G)+\widetilde{\eta}(\mathcal{E}_2)-\widetilde{\eta}(\mathcal{E}_1)\\ &=\operatorname{CS}(\nabla^{\operatorname{ind}^{\mathrm{an}}(E_1)}\oplus\nabla^{\mathrm{flat}},\nabla^{\operatorname{ind}^{\mathrm{an}}(E_2)}\oplus\nabla^{\mathrm{flat}})\\ &=\operatorname{CS}(\nabla^{\operatorname{ind}^{\mathrm{an}}(E_1)}\oplus\nabla^{\mathrm{flat}},\nabla^{\operatorname{ind}^{\mathrm{an}}(E_2)}\oplus\nabla^{\mathrm{flat}})+\operatorname{CS}(\nabla^{\operatorname{ind}^{\mathrm{an}}(G)},\nabla^{\operatorname{ind}^{\mathrm{an}}(G)})\\ &=\operatorname{CS}(\nabla^{\operatorname{ind}^{\mathrm{an}}(E_1)}\oplus\nabla^{\mathrm{flat}}\oplus\nabla^{\operatorname{ind}^{\mathrm{an}}(G)},\nabla^{\operatorname{ind}^{\mathrm{an}}(E_2)}\oplus\nabla^{\mathrm{flat}}\oplus\nabla^{\operatorname{ind}^{\mathrm{an}}(G)})\\ &=\operatorname{CS}(\nabla^{\operatorname{ind}^{\mathrm{an}}(E_1)}\oplus\nabla^{\operatorname{ind}^{\mathrm{an}}(G)},\nabla^{\operatorname{ind}^{\mathrm{an}}(E_2)}\oplus\nabla^{\operatorname{ind}^{\mathrm{an}}(G)})+\operatorname{CS}(\nabla^{\mathrm{flat}},\nabla^{\mathrm{flat}})\\ &=\operatorname{CS}(\nabla^{\operatorname{ind}^{\mathrm{an}}(E_1)}\oplus\nabla^{\operatorname{ind}^{\mathrm{an}}(G)},\nabla^{\operatorname{ind}^{\mathrm{an}}(E_2)}\oplus\nabla^{\operatorname{ind}^{\mathrm{an}}(G)}),\end{aligned}$$

where all the equalities hold up to exact forms. By the variational formula of the Bismut-Cheeger eta form

$$\begin{aligned}\widetilde{\eta}(\mathcal{E}_2)-\widetilde{\eta}(\mathcal{E}_1) &= \mathrm{CS}(\nabla^{\mathrm{ind}^{\mathrm{an}}(E_1)}\oplus\nabla^{\mathrm{ind}^{\mathrm{an}}(G)},\nabla^{\mathrm{ind}^{\mathrm{an}}(E_2)}\oplus\nabla^{\mathrm{ind}^{\mathrm{an}}(G)})\\ &\quad+\int_{X/B}\mathrm{Todd}(\widehat{\nabla}^{T^VX})\wedge\mathrm{CS}(\nabla^2\oplus\nabla^G,\nabla^1\oplus\nabla^G)\mod\Omega^{\mathrm{odd}}_{\mathrm{exact}}(B),\end{aligned}$$

it follows that $\omega\in\Omega^{\mathrm{odd}}_{\mathrm{exact}}(B)$, and therefore $\alpha([\omega])=\alpha(0)=0$. Thus $\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}(\mathcal{E}_1)=\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}(\mathcal{E}_2)$. □

4.2. **The $\mathbb{R}/\mathbb{Z}$ Grothendieck-Riemann-Roch theorem.** In this subsection we give the details of a short proof of the $\mathbb{R}/\mathbb{Z}$ Grothendieck-Riemann-Roch theorem. We first recall a result of Bismut. In the setup of §3.2, with the fibers Spin and $\mathrm{ind}^{\mathrm{an}}(E)\to B$ assumed to form a superbundle, we have

$$\widehat{\mathrm{ch}}(\mathrm{ind}^{\mathrm{an}}(E),\nabla^{\mathrm{ind}^{\mathrm{an}}(E)})+i_2(\widetilde{\eta}(\mathcal{E}))=\widehat{\int_{X/B}}\widehat{\widehat{A}}(T^VX,\nabla^{T^VX})*\widehat{\mathrm{ch}}(E,\nabla^E)\quad(12)$$

[3, Theorem 1.15]. If the fibers are only Spin^c, (7) has the obvious modification

$$\widehat{\mathrm{ch}}(\mathrm{ind}^{\mathrm{an}}(E),\nabla^{\mathrm{ind}^{\mathrm{an}}(E)})+i_2(\widetilde{\eta}(\mathcal{E}))=\widehat{\int_{X/B}}\widehat{\mathrm{Todd}}(T^VX,\widehat{\nabla}^{T^VX})*\widehat{\mathrm{ch}}(E,\nabla^E),\quad(13)$$

for $\widehat{\mathrm{Todd}}(T^VX,\widehat{\nabla}^{T^VX})\in\widehat{H}^{\mathrm{even}}(X;\mathbb{R}/\mathbb{Q})$ the differential character associated to the Todd class as in (4). We will write $\widehat{\mathrm{Todd}}(T^VX,\widehat{\nabla}^{T^VX})$ as $\widehat{\mathrm{Todd}}(\widehat{\nabla}^{T^VX})$ in the sequel. Note that (12) and (13) extend to the general case where $\mathrm{ind}^{\mathrm{an}}(E)\to B$ does not form a bundle [3, p. 23].

The proof of the following proposition essentially uses the same techniques in [8, Proposition 3].

Theorem 1. Let $\pi: X\to B$ be a proper submersion with closed Spin^c fibers of even relative dimension. The following diagram is commutative:

$$\begin{tikzcd} K^{-1}_{\mathrm{L}}(X;\mathbb{R}/\mathbb{Z}) \arrow[r, "\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}"] \arrow[d, "\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}"'] & \widehat{H}^{\mathrm{odd}}(X;\mathbb{R}/\mathbb{Q}) \arrow[d, "\int_{X/B}\mathrm{Todd}(\widehat{\nabla}^{T^VX})\cup(\cdot)"] \\ K^{-1}_{\mathrm{L}}(B;\mathbb{R}/\mathbb{Z}) \arrow[r, "\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}"'] & \widehat{H}^{\mathrm{odd}}(B;\mathbb{R}/\mathbb{Q}) \end{tikzcd}$$

i.e., for a $\mathbb{Z}_2$-graded cocycle $\mathcal{E}=(E^{\pm},h^{\pm},\nabla^{\pm},\phi)\in K^{-1}_{\mathrm{L}}(X;\mathbb{R}/\mathbb{Z})$, we have

$$\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}(\mathcal{E}))=\int_{X/B}\mathrm{Todd}(\widehat{\nabla}^{T^VX})\cup\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathcal{E}).$$

Proof. Consider the difference of differential forms:

$$\begin{aligned}
&\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}(\mathcal{E})) - \int_{X/B} \mathrm{Todd}(\widehat{\nabla}^{T^V X}) \wedge \mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathcal{E}) \\
&= \frac{1}{k}\,\mathrm{CS}(k\nabla^{\mathrm{ind}^{\mathrm{an}}(E)^+}, j^* k\nabla^{\mathrm{ind}^{\mathrm{an}}(E)^-}) + \int_{X/B} \mathrm{Todd}(\widehat{\nabla}^{T^V X}) \wedge \phi \\
&\qquad + \widetilde{\eta}(\mathcal{E}^+) - \widetilde{\eta}(\mathcal{E}^-) - \int_{X/B} \mathrm{Todd}(\widehat{\nabla}^{T^V X}) \wedge \left(\frac{1}{k}\,\mathrm{CS}(k\nabla^{E^+}, j^* k\nabla^{E^-}) + \omega\right) \\
&= \frac{1}{k}\Bigg(\mathrm{CS}(k\nabla^{\mathrm{ind}^{\mathrm{an}}(E)^+}, j^* k\nabla^{\mathrm{ind}^{\mathrm{an}}(E)^-}) \\
&\qquad - \int_{X/B} \mathrm{Todd}(\widehat{\nabla}^{T^V X}) \wedge \mathrm{CS}(k\nabla^{E^+}, j^* k\nabla^{E^-})\Bigg) + \widetilde{\eta}(\mathcal{E}^+) - \widetilde{\eta}(\mathcal{E}^-).
\end{aligned} \tag{14}$$

Here we have chosen the integer k in the Chern-Simon form and the isometric isomorphism j in $\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}(\mathcal{E}))$ and in $\int_{X/B} \mathrm{Todd}(\widehat{\nabla}^{T^V X}) \wedge \mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathcal{E})$ to be the same. This is valid since the cohomology classes of these odd forms do not depend on the choices made.

Applying the map i_2 (3) to (14), we get

$$\begin{aligned}
&i_2\Bigg(\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}(\mathcal{E})) - \int_{X/B} \mathrm{Todd}(\widehat{\nabla}^{T^V X}) \wedge \mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathcal{E})\Bigg) \\
&= \widehat{\mathrm{ch}}(\mathrm{ind}^{\mathrm{an}}(E)^+, \nabla^{\mathrm{ind}^{\mathrm{an}}(E)^+}) - \widehat{\mathrm{ch}}(\mathrm{ind}^{\mathrm{an}}(E)^-, \nabla^{\mathrm{ind}^{\mathrm{an}}(E)^-}) \\
&\qquad - \widehat{\int_{X/B}} \widehat{\mathrm{Todd}}(\widehat{\nabla}^{T^V X}) * (\widehat{\mathrm{ch}}(E^+, \nabla^{E^+}) - \widehat{\mathrm{ch}}(E^-, \nabla^{E^-})) + i_2(\widetilde{\eta}(\mathcal{E}^+)) \\
&\qquad - i_2(\widetilde{\eta}(\mathcal{E}^-))
\end{aligned}$$

by 5 of Proposition (1) and 4 of Proposition (2). Since

$$\begin{aligned}
&\widehat{\mathrm{ch}}(\mathrm{ind}^{\mathrm{an}}(E)^+, \nabla^{\mathrm{ind}^{\mathrm{an}}(E)^+}) - \widehat{\mathrm{ch}}(\mathrm{ind}^{\mathrm{an}}(E)^-, \nabla^{\mathrm{ind}^{\mathrm{an}}(E)^-}) \\
&= \widehat{\mathrm{ch}}(\ker(\mathsf{D}^{E^+})^+, \nabla^{\ker(\mathsf{D}^{E^+})^+}) + \widehat{\mathrm{ch}}(\ker(\mathsf{D}^{E^-})^-, \nabla^{\ker(\mathsf{D}^{E^-})^-}) \\
&\qquad - \widehat{\mathrm{ch}}(\ker(\mathsf{D}^{E^+})^-, \ker(\mathsf{D}^{E^+})^-) + \widehat{\mathrm{ch}}(\ker(\mathsf{D}^{E^-})^+, \ker(\mathsf{D}^{E^-})^+) \\
&= \widehat{\mathrm{ch}}(\mathrm{ind}^{\mathrm{an}}(E^+), \nabla^{\mathrm{ind}^{\mathrm{an}}(E^+)}) - \widehat{\mathrm{ch}}(\mathrm{ind}^{\mathrm{an}}(E^-), \nabla^{\mathrm{ind}^{\mathrm{an}}(E^-)}),
\end{aligned}$$

it follows that (14) becomes

$$\begin{aligned}
&\widehat{\mathrm{ch}}(\mathrm{ind}^{\mathrm{an}}(E^+), \nabla^{\mathrm{ind}^{\mathrm{an}}(E^+)}) - \widehat{\mathrm{ch}}(\mathrm{ind}^{\mathrm{an}}(E^-), \nabla^{\mathrm{ind}^{\mathrm{an}}(E^-)}) \\
&- \widehat{\int_{X/B}} \widehat{\mathrm{Todd}}(\widehat{\nabla}^{T^V X}) * (\widehat{\mathrm{ch}}(E^+, \nabla^{E^+}) - \widehat{\mathrm{ch}}(E^-, \nabla^{E^-})) + i_2(\widetilde{\eta}(\mathcal{E}^+)) \\
&\qquad - i_2(\widetilde{\eta}(\mathcal{E}^-)) \\
&= 0
\end{aligned}$$

by (13). It follows from the definition of the map i_2 that

$$\mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathrm{ind}^{\mathrm{an}}_{\mathrm{L}}(\mathcal{E})) - \int_{X/B} \mathrm{Todd}(\widehat{\nabla}^{T^V X}) \wedge \mathrm{ch}_{\mathbb{R}/\mathbb{Q}}(\mathcal{E}) \in \Omega^{\mathrm{odd}}_{\mathbb{Q}}(B). \tag{15}$$

The theorem is proved by reducing (15) by $\mathbb{Q}$ and taking the cohomology class by regarding (15) as a cocycle. □

REFERENCES

1. M.F. Atiyah, V. K. Patodi, and I.M. Singer, *Spectral asymmetry and Riemannian geometry II*, Math. Proc. Cambridge Philos. Soc. **78** (1975), 405–432.
2. N. Berline, E. Getzler, and M. Vergne, *Heat kernels and Dirac operators*, Grundlehren Text Editions, Springer-Verlag, Berlin, 2004.
3. J.M. Bismut, *Eta invariants, differential characters and flat vector bundles*, Chinese Ann. Math. Ser. B **26** (2005), 15–44.
4. J.M. Bismut and J. Cheeger, *η-invariants and their adiabatic limits*, J. Amer. Math. Soc. **2** (1989), 33–70.
5. J. Cheeger and J. Simons, *Differential characters and geometric invariants*, in Geometry and Topology (College Park, Md., 1983/84), Lecture Notes in Math. **1167** (1985), 50–80.
6. X. Dai, *Adiabatic limits, nonmultiplicativity of signature, and Leray spectral sequence*, J. Amer. Math. Soc. **4** (1991), 265–321.
7. D. Freed and J. Lott, *An index theorem in differential K-theory*, Geom. Topol. **14** (2010), 903–966.
8. M.H. Ho, *A condensed proof of the differential Grothendieck-Riemann-Roch theorem*, arXiv:1111.5546v5, To appear in the Proc. Amer. Math. Soc.
9. M. Hopkins and I.M. Singer, *Quadratic functions in geometry, topology,and M-theory*, J. Diff. Geom. **70** (2005), 329–425.
10. M. Karoubi, *Homologie cyclique et K-théorie*, Astérisque **149** (1987).
11. J. Lott, *$\mathbb{R}/\mathbb{Z}$ index theory*, Comm. Anal. Geom. **2** (1994), 279–311.

DEPARTMENT OF MATHEMATICS, BOSTON UNIVERSITY
E-mail address: `homanho@bu.edu`